

GENERAL INDEX TO VOLUME XLII

New scientific names of plants and the final members of new combinations are printed in bold face type; synonyms and page numbers having reference to figures and plates, in *italics*; and all other matter in ordinary type.

A

- Acanthothamnus* *aphyllus*, 285
Acarospora (*Pachnolepia*) *Gwynnii*, 144
Adélie Land, lichen flora of, 143
Adlay, 105; chromosome morphology of, 115, 127, 128
Alectoria *congesta*, 143, 145
Allosorus *andromedaefolius*, 102
Alsodecia *glabra*, 231
Anaxagorea *Allenii*, 151; *crassipetala*, 151; *dolichocarpa*, 151
 Animal products, as a source of food in Karimpur, 309, 344
Annonaceae, Three new, from Panama, 151
 Antarctic continent and the subantarctic islands, Lichenological notes on the flora of the, I-IV, 131
Anthoxanthum, 113
Apluda *digitata*, 106
Argopsis *cymosa*, 134
 Asiatic relatives of maize, Studies on, 103
Aspicilia *disjuguenda*, 135; *lygemma*, 135
Aspiciliopsis *macrophthalma*, 135
 Australian National Antarctic Research Expedition, lichens collected by, 131

B

- Balansia* *trinitensis*, 136
Blastenia *keroplasta*, 135
 "Blue Adlay", 107, 108, 109; chromosome morphology of, 115, 127, 128
 Botanical catalogues of Auguste de St. Hilaire, 153
Brahmans, food habits of, 307, 379, 400
 Brazil, botanical collectors in, 153
Briza, 113
Buellia *frigida*, 143, 146; *Johnstoni*, 146; *Mawsoni*, 143; *subplicata*, 135

C

- Caloplaca* *crozetica*, 135
 Catalogues, botanical, of Auguste de St. Hilaire, 153
Catba *bentharii*, 244; *monosperma*, 249; *vitiensis*, 291

- Catillaria* (*Eucatillaria*) *Rudolphii*, 139
Ceanothus *paniculatus*, 230
 Cedar, Red, see *Juniperus virginiana*
Celastrus, A revision of the genus, 215
Celastrus, 227; *aculeatus*, 274; *acuminatus*, 285; *adenophylla*, 285; *alatus*, 285; *alpestris*, 285; *angulatus*, 239; *annamensis*, 244; *aphyllus*, 285; *apoensis*, 246; *approximata*, 250; *aquifolius*, 285; *articulata* var. *cuneata*, 261, var. *orbiculata*, 260; *articulatus*, 260, var. *pubescens*, 261, var. *papillosus*, 261, var. *punctatus*, 266, var. *stephanotiifolius*, 261; *attenuatus*, 285; *australis*, 236; *axillaris*, 250; *bentharii*, 244; *bilocularis*, 285; *bivalvis*, 285; *boaria*, 285; *bodinieri*, 285; *bullatus*, 240; *buxifolia*, 285; *cantonensis*, 249; *casearii-folius*, 282; *cavaleriei*, 265, 286; *championii*, 244, 246; *chiapensis*, 282; *chungii*, 286; *ciliidens*, 276; *circumcissus*, 286; *colombianus*, 280; *clemacanthus*, 276; *confertus*, 286; *crassifolia*, 273; *crenatus*, 286; *crenulatus*, 286; *crispulus*, 266; *cuneifolius*, 286; *cunninghamii*, 286, var. *parviflora*, 286; *dependens*, 231; *dilatatus*, 286; *discolor*, 285; *disperma*, 286; *diversifolia*, 286; *dubia*, 286; *elevativenus*, 267; *emarginatus*, 286; *esquirolianus*, 286; *esquirolii*, 287; *euonymoidea*, 287; *euphlebiphyllus*, 231; *fasciculatus*, 287; *feddei*, 287; *finlaysonianus*, 287; *flagellaris*, 276; *floribundus*, 287; *fournieri*, 287; *geminiflorus*, 267; *gemma*, 258; *gemma*, 258; *glaucophyllus*, 253; *glaucus*, 287; *gracilimus*, 267; *grenadensis*, 285; *haenkea*, 287; *hamelii*, 287; *heterophyllus*, 287; *heyneana*, 287; *hindsii*, 249, var. *henryi*, 249; *hirsutus*, 270; *hookeri*, 254; *hookeri*, 274; *hypoglaucus*, 257; *hypoleucus*, 257, f. *β*. *argutior*, 257, f. *α*. *genuina*, 257, f. *γ*. *puberula*, 273; *ilicifolius*, 287; *insularis*, 261; *japonicus*, 287; *jeholensis*, 261; *jodinii*, 287; *kiusianus*, 266; *kouytschensis*, 287; *kusanoi*, 268; *lancifolia*, 261; *laotica*, 231; *latifolius*, 239; *leiocarpus*, 267; *lenticellatus*, 278; *leptopus*, 287; *liebmannii*, 279; *linearis* var. *madagascariensis*, 287;

- lineatus*, 287; *listeri*, 288; *loeseneri*, 265; *lokchongensis*, 258; *longe-racemosus*, 267; *longipes*, 281; *lucida*, 291; *lucida*, 288; *lycioides*, 288; *lyi*, 288; *macrocarpus*, 288; *madagascariensis*, 238; *magellanicus*, 288; *mainisiana*, 279; *mairei*, 288; *malayensis*, 246; *marianensis*, 250; *mauritanica*, 288; *maytenus*, 288; *membranifolius*, 255; *meridensis*, 284; *merrillii*, 250; *metzianus*, 230; *mexicanus*, 288; *micrantha*, 288; *microcarpus*, 285; *mollis*, 288; *monosperma*, 249; *monospermoides*, 246; *monospermus*, 244; *montana*, 288; *moya*, 288; *muelleri*, 288; *multiflorus*, 231; *myrsiflorus*, 288; *neglecta*, 272, 288; *nepalensis*, 288; *novoguineensis*, 234; *nutans*, 229; *oblanceifolia*, 274; *oblongifolius*, 250; *obtusatus*, 288; *obtusifolia*, 288; *octogonus*, 288; *opposita*, 289; *orbiculata* f. *major* 261, f. *microphylla*, 261, var. *aureo-arillata*, 261; *orbiculatus*, 260; *orbiculatus*, 253, f. *papillosus*, 261, var. *papillosus*, 261, var. *pilosus*, 261, var. *punctatus*, 266; *orixa*, 289; *ovalifolius*, 289; *ovatus*, 289; *oxyphyllus*, 289; *pachyrachis*, 279; *pallidus*, 289; *panamensis*, 280; *paniculatus*, 229, var. *andamanica*, 230, var. *Balansae*, 236, ssp. *multiflorus*, 231, ssp. *paniculatus*, 229, ssp. *serratus*, 231, var. *pubescens*, 230; *papuana*, 236; *parviflorus*, 289; *parvifolius*, 289; *pauciflora*, 289; *pentagyna*, 289; *polybotrys*, 231; *pringlei*, 281; *pubescens*, 230; *punctatus*, 266; *punctatus*, 261, var. *microphyllus*, 267; *punctulatus*, 266; *quadrangulatus*, 289; *racemosus*, 279, var. *trinitensis*, 285; *racemulosus*, 249; *repandus*, 285; *reticulatus*, 285; *retusa*, 289; *rhombifolius*, 289; *richardi*, 289, γ. *latifolius*, 289, δ. *coclearifolius*, 289, ε. *elaeodendroides*, 289; *richii*, 237; *rigida*, 289; *robustus*, 289; *rosthornianus*, 265; *rothiana*, 229; *rothianus*, 290; *royleanus*, 290; *rufa*, 290; *rugosus*, 253; *salicifolia*, 290; *scandens*, 240; *scandens*, 266; *seguini*, 290; *semiarillata*, 290; *sepiarius*, 290; *serrulatus*, 290; *siltepecanus*, 282; *spicatus*, 290; *spiciformis*, 256, var. *laevis*, 256; *spinifolius*, 290; *spinosus*, 290; *stephanotiifolius*, 261; *striatus*, 266, 290; *strigillosus*, 261; *stylosa*, 254, 263, var. *Loeseneri*, 254; *stylosus*, 272, ssp. *stylosus*, 272, ssp. *glaber*, 273; *suaveolens*, 290; *subspicatus*, 236; *tatarinowii*, 260; *tetramerus*, 290; *tristis*, 290; *uncinatus*, 290; *vanioti*, 256; *variabilis*, 290; *venulosus*, 249; *versicolor*, 261; *verticillata*, 290; *verticillatus*, 290; *vitiensis*, 291; *vulcanicofolus*, 282; *wallichianus*, 291; *wallichii*, 291; *wightianus*, 291; *yunnanensis*, 291; *zeylanica*, 291
- Cenomyce phyllophora*, 134; *sarmentosa*, 140
- Cereals used in North India, 329
- Chaillietia gelonioides*, 285
- Chionachne*, 103, 106, 108, chromosome morphology of, 114, 125, 130; *biaurita*, 106; *Hubbardiana*, 106; *Koenigii*, 106, 107, 111, 112, 113, chromosome morphology of, 117, 125, 130; *Massii*, 106; *semiteres*, 106
- Chlorocypbella*, 136; *subtropica*, 136
- Chromosome morphology of *Coix*, *Polytoca* and *Chionachne*, 114
- Chromosome numbers: in *Celastrus*, 225; in *Coix*, 113, in *Selaginella*, 11
- Cladia aggregata*, 140
- Cladonia floriformis*, 140; *Johnstoni*, 134; *Mawsoni*, 140; *phyllophora*, 134; *sarmentosa*, 140
- Coenogonium subtorulosum*, 137
- Coix*: cytological study of, 107, materials used, 107, methods, 113; history of, 103; nomenclature in the genus, 104
- Coix*, 103; *agrestis*, 104; *aquatica*, 104, 108, 113; *arundinacea*, 104, 106; *barbata*, 106; *chinensis*, 104; from Cuttack, India, 108; *cryptoides*, 106; *exaltata*, 104; *gigantea*, 104, 109, 110, 116, 125, 128, 129, f. *aquatica*, 104, 114; *heteroclita*, 106; *Koenigii*, 106; *Lachryma*, 104, var. *stenocarpa*, 113; *Lachryma-Jobi*, 104, 108, f. *agrestis*, 104, var. *agrestis*, 104, f. *aquatica*, 104, var. *aquatica*, 104, var. *gigantea*, 104, var. *major*, 104, f. *aquatica*, 104, f. "mayuen", 104, var. "Mayuen", 104, 113, var. *Ma-yuen* (1), 116, 124, var. *Ma-yuen* (2), 116, 124-128, var. *minor*, 104, f. *monilifer*, 103, var. *monilifer*, 104, f. *palustris*, 104, var. *palustris*, 104, f. *stenocarpa*, 104, var. *stenocarpa*, 104, var. *typica* "Blue Adlay", 105, 108, 115, 124, 126-128; *lingulata*, 104; *Ma-yuen*, 104; *ouwehandii*, 104; *palustris*, 104; *pendula*, 104; *poilanei*, 104, 108, 109, 113, 114, 116, 124-128; *puellarum*, 104; *stenocarpa*, 104; *tubulosa*, 104; from western Ghats, India, 109
- Collemopsisidium pyrenuloides*, 133
- Colubrina ferruginosa*, 289
- Coyx tubulosa*, 104
- Cuvier's account of St. Hilaire's travels, 154
- Cyathorhachis Wallichiana*, 106
- Cyphella aeruginascens*, 136
- Cytology of *Coix* and its relatives, 115

D

- "Dal Roti" in India, 333
 Dale, Ernest E., Norton H. Nickerson and:
 Tassel modifications in *Zea Mays*, 195
Deuterolichenes, 136
Diosma serrata, 231
Doryopteris, a new species of, from Suri-
 nam, 213
Doryopteris, 213; *collina*, 214; *conformis*,
 213, 214; *lomariacea*, 213; *pedata* var.
multipartita, 214; *sagittifolia*, 214; *vari-*
ans, 214
 Dreuzy's appendix to St. Hilaire's "Voyage
 à Rio Grande do Sol," 157
 Dwyer, John D.: The botanical catalogues
 of Auguste de St. Hilaire, 153

E

- Ear of *Zea Mays*, morphology of, 195
 Ecological factors in relation to distribution
 of *Juniperus*, 187
Elaeodendron glaucum, 287
Eleonurus candidus var. *bisetosus*, 167
Embelia esquirolii, 258
Ephelidium, 136; *heardense*, 136
Ephelis, 136; *trinitensis*, 136
Erythrospermum hypoleucum, 257
Euonymus altus, 290; *cuneifolius*, 286;
euphlebiophyllus, 231; *thunbergianus*, 285
Euonymoides, 227; *scandens*, 240

F

- Ferns, See *Pellaea* and *Selaginella*
 Festival dishes of North India, 346
Flueggia serrata, 249
 Food: Coix as, 105; of North India, 303;
 diet of Brahmans, 307, 379, of other
 castes, 381, 401
 Foods of a Hindu Village in North India,
 303
 Forage plants, 103
 Frequency distribution of three species of
Celastrus, 232
 Fries, Robert E.: Three new *Annonaceae*
 from Panama, 151
 Fruits eaten in India, 316, 341

G

- Gasparrinia Harrissoni*, 146; *macquariensis*,
 142
 Geographical distribution of Coix, relation-
 ship between chromosome number and,
 120
 "Ghi" as food in North India, 368
 Glade in Missouri Botanical Garden Arbo-
 retum, 173, 190; species list with fre-
 quencies for under story of, 178

- Gouania corylifolia*, 290
Gramineae, 103
 Gray Summit, Mo., history of land usage
 at, 173
 Greens used as food in India, 341
Grewia feddei, 287; *henryi*, 287
Guatteria inuncta var. *caudata*, 152
Gymnosporia acuminata, 289; *crenata*, 286;
diversifolia, 286; *emarginata*, 286; *heyne-*
ana, 287; *montana*, 286, 288, 289; *neg-*
lecta, 285, 288; *neglecta*, 272; *ovata*,
 289; *rothiana*, 290; *royleana*, 290; *rufa*,
 290; *trigyna*, 288; *variabilis*, 290; *viti-*
ensis, 286, 291; *wallichiana*, 286, 289,
 291

H

- Hall, Marion Trufant: Comparison of Juni-
 per populations on an Ozark Glade and
 Old Fields, 171, Appendix, 192
 Heard Island: Additions to the Lichen flora
 of, 131; views of, 149
Heppia antarctica, 144
 Heredity, *Zea Mays* as a tool in study of,
 195
 Hindu village in North India, foods of a,
 303
 Hou, Ding: A revision of the genus *Celas-*
trus, 215
 Hybridization in *Celastrus*, 225

I

- Ilex crenata*, 285; *cuneifolia*, 287; *macro-*
carpa, 290; *purpurea*, 285; *scandens*, 286;
suaveolens, 290
 India: Coix from, 108, 109; Foods of a
 Hindu village in, 303; food plants of, 103
 Inflorescence of *Zea Mays*: staminate, see
 Tassel; pistillate, see Ear
 Inflorescences of *Celastrus*, 219; diagrams
 of, 220, 221
 Introgression in *Juniperus*, 171, 186
Iodina rhombifolia, 289

J

- Job's Tears, see Coix
 Juniper populations on an Ozark glade and
 old fields, Comparison of, 171
Juniperus, 171; *Ashei*, 171, variation in,
 181, 182; *horizontalis*, 171; *virginiana*,
 171, variation in, 181, 183, Eastern race,
 171, Northern race, 171, 186, Ozark
 race, 171, 186; *virginiana* var. *crebra*,
 171, 186, var. *typica*, 171; *virginiana* ×
J. Ashei, 186
Juniperus virginiana and *J. Ashei*, variation
 patterns in, 181, 182, 184, 185

K

- Karimpur, India, foods of, 303
 Kramer, Karl U., and Rolla M. Tryon, Jr.:
 A new species of *Doryopteris* from Suri-
 nam, 213
 Kurrima sp., 289
 Kuttlingeria crozetica, 135

L

- Leaves of *Juniperus*, variations in, 181, 182,
 184, 185
Lecanora chrysouleuca v. *melanophthalma* f.
exsulans, 145; *dichroa*, 134; *disjunctuenda*,
 135; *exsulans*, 145; *griseomarginata*, 145;
perrugosa, 140; *versicolor*, 137
Lecidea assentiensis, 134; Auberti, 134;
 Harrissoni, 144; *lygomma*, 135; sub-
 assentiensis, 134; subglobulata, 139; sub-
 plicata, 135
Lichen aggregatus, 140
 Lichenological notes on the flora of the
 Antarctic continent and the subantarctic
 islands, I-IV, 131
Lichina antarctica, 132
Lopadium perpallidum, 136; Deightoni, 136
Lycopodium bryoides, 40, 67; *carinatum*,
 82; *ciliatum*, 77; *Dregei*, 57; *laxum*, 82;
rupestris, 64; *struthioloides*, 81, 82

M

- Macquarie Island, additions to the Lichen
 flora of, 137
 Mac Robertson Land, additions to the flora
 of, 143
 Maize: in North India, 329; Studies on
 Asiatic relatives of, 103; Tassel modifica-
 tions in, 195
 Mangoes in North India, 316
Maytenus bilocularis, 285; *boaria*, 285, 288,
 290; *buxifolius*, 289; *cochlearifolius*, 289;
confertus, 286; *cunninghamii*, 286; *dis-*
permus, 286; *elaeodendroides*, 289;
fournieri, 287; *lineatus*, 287; *macro-*
carpus, 288; *magellanicus*, 288; *meriden-*
sii, 284; *orbicularis*, 286, 288, 289; *ovata*,
 289; *quadrangulatus*, 289; *racemosus*,
 279; *retusa*, 286, 289; *truncatus*, 287;
verticillatus, 290; *williamsii*, 282
 Meiosis in *Coix* and related genera, 118
Menegazzia circumscoriata, 142
Microglaua kerguelena, 132
Microthelia macquariensis, 137
Microtropis bivalvis, 285
 Missouri Botanical Garden Arboretum, 172:
 Junipers at Cedar Hill, 175; The Glade,
 173; The Old Field, 174; species list with
 frequencies for understory of Glade and
 Old Field, 178

Morphology of tassel of *Zea Mays*, 195

Moya spinosa, 288

- Mutants of tassel of *Zea Mays*: descriptions
 of, 199; key to, 196; measurements of
 morphological features, 197; spikelet pairs
 on, 211; Branched Silkless, pl. 28; Club,
 pl. 27; male-sterile, pl. 27; *Ramosa* 1
 and *Ramosa* 2, pl. 29; Standard, pl. 22;
 Tassel-seed 1 and Tassel-seed 2, pl. 23;
 Tassel-seed 3 and Tassel-seed 4, pl. 24;
 Tassel-seed 5 and Tassel-seed 6, pl. 25;
 Tassel-seed 7 and Tassel-seed 8, pl. 26;
 Vestigial glume, pl. 28

Myrsine semiserrata, 286, 290

N

- Neurospora aeromelanus* v. *inactivus* f.
scabridulus, 143; *insularis*, 135; *trachy-*
carpus, 135
 Nickerson, Norton H., and Ernest E. Dale:
 Tassel modifications in *Zea Mays*, 195
 Nirodi, Nalini: Studies on Asiatic relatives
 of maize, 103
Nostoc, sp., 146
 Nutrition practices in a Hindu village of
 North India, 356

O

- Old fields in Missouri Botanical Garden
 Arboretum, 174, 190, 191; distribution
 of Juniper seedlings in, 180; variations
 of junipers in, 180, 181; species list with
 frequencies for understory of, 178
Orixa japonica, 286, 287, 289
 Ozark glade and old fields, Comparison of
 Juniper populations on, 171

P

- Pannaria dichroa*, 134; *glaucella*, 132
Parmelia Johnstoni, 145; *pubescens* v. *con-*
gesta, 145; *sublugubris*, 142; *tenuirima*,
 142; *turgidula*, 142
 Pellaea, A new, from South Africa, 101
Pellaea andromedaefolia, 102; *intermedia*,
 102; *ovata*, 102; *myrtillifolia*, 101; *rufa*,
 101, 102
Peltigera Lairdi, 138
 People of India, 304
Perrottetia alpestris, 285
Pertusaria tyloplaca, 140
Phyllanthus sp., 290
Pittosporum floribundum, 288, 290
Placodium bicolor, 135
Placopsis bicolor, 135; *perrugosa*, 140
Pleurostelia cochinchinensis, 289, 291
Polytoca, 103, 108; *barbata*, 113; *bracteata*,
 106; *cyathopoda*, 106; *digitata*, 106;
javanica, 106; *macrophylla*, 106, 107,

- 110, 111, 113, chromosome morphology of, 117, 126, 129, 130; Massii, 106; semiteres, 106; Wallichiana, 106
 Populations, Juniper, comparisons of, on an Ozark glade and old fields, 171; Appendix, 192
Prema parviliba, 291
 Preservation of food in North India, 323
Protoplastenia citrina, 145
Prunus myrtifolia, 288
Pseudocypbellaria glabra, 138
Psoroma versicolor, 137
 Pulses, use of, as food in North India, 333, 337

R

- Ramalina geniculata*, 142
 Red Cedar, see *Juniperus virginiana*
Rhamnus crenatus, 286, 287; *esquirolii*, 288; *napalensis*, 290; *racemosus*, 287
Rhizocarpon Johnstoni, 134; *keruelense*, 134; *Mawsonii*, 134
Rinodina aspicilina, 136; *frigida*, 143, 146
 Rudolph, Emanuel D., Carroll W. Dodge and: Lichenological notes on the flora of the Antarctic continent and the subantarctic islands, I-IV, 131

S

- Sabia parviflora* var. *nitidissima*, 287; *yunnanensis*, 288
 St. Hilaire, Auguste de, The botanical catalogues of, 153; numbers of collections listed under specific families, 159; numerical sequences of, 158; page of Catalogue C², 168; page from list of collection numbers, 170; role of catalogues in taxonomic monographs, 163; Weddell's key to, 161, 169
Saurauja vaniata, 256
Schoepfia flexuosa, 287
Sclerachne, 103; *cyathopoda*, 106; *punctata*, 113
Scutia commersonii, 291; *paniculata*, 230
Selaginella rupestris and its allies, 1
Selaginella, 16; *acanthonota*, 26; *amazonica*, 34; *Arachavaletae*, 34; *arenaria*, 26; *arenicola*, 23, ssp. *acanthonota*, 26, 94, ssp. *arenicola*, 26, 94, ssp. *Riddellii*, 24; *arizonica*, 78, 99; *Arsenei*, 41, 95; *Aschenbornii*, 77; *asprella*, 75, 98; *Balansae*, 27, 94; *Bigelovii*, 22, 94; *Bolanderi*, 49; *Bourgeauii*, 64; *bryoides*, 40; *caffrorum*, 53, 96; *capensis*, 53; *carinata*, 50, 96; *Chrismarii*, 20, var. *Karwinskyana*, 20, var. *Necana*, 20; *cinerascens*, 40, 95; *columbiana*, 67; *Coryi*, 28; *deflexa*, 16; *densa*,

- 66, var. *densa*, 66, 97, 98, var. *scopulorum*, 97, var. *Standleyi*, 71, 97; *denticulata*, 11; *Dregei*, 57, 96, var. *Bachmanniana*, 57, var. *Hildebrandtiana*, 57, var. *Petersiana*, 57, var. *pretoriensis*, 57, var. *Rehmanniana*, 57, var. *Welwitschiana*, 57; *echinata*, 54, 96; *Engelmannii*, 68, *eremophila*, 80, 99; *extensa*, 47, 95; *Fendleri*, 62; *floridana*, 26; *funiformis*, 26; *gracillima*, 17; *grisea*, 57; *Hansenii*, 49, 96; *Haydenii*, 76; *helvetica*, 11; *Hintonii*, 37; *humifusa*, 26; *indica*, 52, 96; *Landii*, 81, 99; *lepidophylla*, 8; *leucobryoides*, 74, 98; *longipila*, 52, 68; *macrathera*, 41, 95; *Mildei*, 34; *montanensis*, 43; *montevideensis*, 34; *mutica*, 44, var. *limitanea*, 46, var. *mutica*, 46, 95, var. *texana*, 46; *muticola* × *rupincola*, 21, × *neomexicana*, 21; *nivea*, 55, 96; *njamnamensis*, 52, 96; *oregana*, 61, 97; *Parishii*, 80, 99; *pauciciliata*, 34; *peruviana*, 77, 99, var. *Dombeyana*, 77; *Phillipsiana*, 40; *porrecta*, 37; *Preissiana*, 17; *proxima*, 56, 96; *pumila*, 17; *pygmaea*, 17; *Quartiniana*, 53; *Riddellii*, 24; *Rosendablii*, 43; *rupestris*, 64, 97, f. *amazonica*, 34, f. *amurensis*, 71, f. *Bolanderi*, 49; *Bourgeauii*, 64, f. *brasiliensis*, 34, f. *caffrorum*, 53, f. *Dregei*, 57, f. *Engelmannii*, 70; *Hansenii*, 49, f. *Haydenii*, 70, f. *indica*, 52, f. *longipila*, 68, f. *manchurienensis*, 71, f. *montanensis*, 43, f. *peruviana*, 77, f. *Schmidtii*, 71, f. *sibirica*, 71, f. *Wallacei*, 43, f. *Wrightii*, 48, ssp. *Balansae*, 27, var. *acanthonota*, 26, var. *Balansae*, 27, var. *Bolanderi*, 49, var. *borealis*, 81, var. *brasiliensis*, 34, var. *brevipila*, 81, var. *columbiana*, 43, var. *densa*, 70, var. *Fendleri*, 62, var. *Hansenii*, 49, var. *incurva*, 53, f. *abyssinica*, 53, f. *angolensis*, 53, f. *capensis*, 53, var. *longipila*, 81, subvar. *glauca*, 81, subvar. *viridis*, 81, var. *mexicana*, 47, 77, var. *montanensis*, 43, var. *peruviana*, 77, var. *recurva*, 57, f. *Dregeana*, 57, f. *Welwitschiana*, 57, var. *rupincola*, 20, var. *Sartorii*, 37, var. *shakotanensis*, 43, var. *Sherwoodii*, 29, var. *tortipila*, 29, var. *tropica*, 82, var. *Wallacei*, 43; *rupincola*, 20, 94; *Sartorii*, 36, 95, var. *oregonensis*, 36, var. *venezuelensis*, 36; *Schmidtii*, 71, var. *Krauseorum*, 71; *scopulorum*, 67; *selaginoides*, 16; *Sellowii*, 34, 95; *shakotanensis*, 43, 95; *Sbeldonii*, 77; *Sherwoodii*, 29; *sibirica*, 71, 98; *spinulosa*, 11; *Standleyi*, 71; *Steyermarkii*, 49, 95; *struthioloides*, 82; *tortipila*, 29, 94; *uliginosa*, 17; *Underwoodii*, 62, 97,

var. *dolicbotricha*, 62; *utahensis*, 74, 98;
Vardei, 61, 97; *viridissima*, 28, 94;
Wallacei, 43, 95, f. *columbiana*, 43;
Watsonii, 72, 98; var. *mutica*, 46; *Weatherbiana*, 28, 94; *Wightii*, 39, var. *Philipsiana*, 40, var. *vetusta*, 39, var. *Wightii*, 39; *Wrightii*, 48, 95; *Wrightii*, 48

Semmarilla, 227

Sempervirentes, 242

Simmondsia californica, 288

Siphulastrum cladinoideis, 133

South Africa, A new *Pellaea* from, 101

Spices used in cooking food in North India, 319

Sporopodium, 136

Stachygynandrum rupestre, 94

Steinera glauca, 132; *Werthii*, 132

Stereocaulon argodes, 140; *Argus*, 140;

cymosum, 134; *pulvinare*, 140; *submollescens*, 140

Stricta glabra, 138

Surinam, A new species of *Doryopteris* from, 213

T

Tassel modifications in *Zea Mays*, 195

Tassels of *Zea Mays*: morphology of, 195; certain mutants of, 196, pls. 22-29, relationships between, and normal tassels, 198

Taxonomic monographs, role of St. Hilaire's catalogues in, 163

Thamnolecania macquariensis, 141

Thelidium heardense, 131; *praevalescens*, 131

Tichorhynchium, 133

Tomatoes, use of, in North India, 309

Trichilia jodini, 287

Trilobachne, 103

Trinidad, Coix "Blue Adlay" from, 108

Tripterogium wilfordii, 286

Tryon, Alice F.: A new *Pellaea* from South Africa, 101

Tryon, Rolla M., Jr.: *Selaginella rupestris* and its allies, 1; Karl U. Kramer and: A new species of *Doryopteris* from Surinam, 213

U

Umbilicaria antarctica v. *subvirginis*, 144; *Hunteri*, 143; *spongiosa* v. *subvirginis*, 144; *subcerebriformis*, 144

United Provinces of India, food of, 304

Unonopsis costaricensis, 152; *guatterioideis*, 152; *obovata*, 152; *panamensis*, 151;

Pittieri, 152; *Schippii*, 152

Urceolaria macrophthalmia, 135

Urecolina kergueliensis, 134

Urostachys carinatus, 82

Usnea antarctica, 143; *arida* v. *muscolola*, 142; *contexta*, 142; *dasyogoides*, 142; *insularis*, 135; *pustulata*, 143; *scabridula*, 143; *Taylori*, 135; *torulosa*, 142; *trachycarpa*, 135; *xanthopoga*, 142

V

Variation in *Juniperus* at the Garden Arboretum, 181

Vegetables grown in North India, 315, 336

Verrucaria kerguelena, 132; *praevalescens*, 131

W

Weddell's key to catalogues of St. Hilaire, 161; portion of page from, 169

Wiser, Charlotte Viall: Foods of a Hindu Village in North India, 303

Wimmeria mexicana, 288

Wheat: grown in India, 330; bread from, 333

X

Xanthopyrenia heardensis, 132

Xanthoria Mawsoni, 143, 146

Z

Zea Mays: Tassel modifications in, 195; tassel mutants, illustrations of, pls. 22-29, key to, 196

955]

44;
nis,

des,
51;

ola,
42;
ula,
ny-

bo-

36
ms,

re,

du

m,

95;
2-